


APMO 1991 – Problems and Solutions

Problem 1

Let G be the centroid of triangle ABC and M be the midpoint of BC . Let X be on AB and Y on AC such that the points X, Y , and G are collinear and XY and BC are parallel. Suppose that XC and GB intersect at Q and YB and GC intersect at P . Show that triangle MPQ is similar to triangle ABC .

Solution 1

Let R be the midpoint of AC ; so BR is a median and contains the centroid G .



It is well known that $\frac{AG}{AM} = \frac{2}{3}$; thus the ratio of the similarity between AXY and ABC is $\frac{2}{3}$. Hence $GX = \frac{1}{2}XY = \frac{1}{3}BC$.


Now look at the similarity between triangles QBC and QGX :

$$\frac{QG}{QB} = \frac{GX}{BC} = \frac{1}{3} \implies QB = 3QG \implies QB = \frac{3}{4}BG = \frac{3}{4} \cdot \frac{2}{3}BR = \frac{1}{2}BR.$$

Finally, since $\frac{BM}{BC} = \frac{BQ}{BR}$, MQ is a midline in BCR . Therefore $MQ = \frac{1}{2}CR = \frac{1}{4}AC$ and $MQ \parallel AC$. Similarly, $MP = \frac{1}{4}AB$ and $MP \parallel AB$. This is sufficient to establish that MPQ and ABC are similar (with similarity ratio $\frac{1}{4}$).

Solution 2

Let S and R be the midpoints of AB and AC , respectively. Since G is the centroid, it lies in the medians BR and CS .



Due to the similarity between triangles QBC and QGX (which is true because $GX \parallel BC$), there is an inverse homothety with center Q and ratio $-\frac{XG}{BC} = \frac{XY}{2BC}$ that takes B to G and C to X . This homothety takes the midpoint M of BC to the midpoint K of GX .

Now consider the homothety that takes B to X and C to G . This new homothety, with ratio $\frac{XY}{2BC}$, also takes M to K . Hence lines BX (which contains side AB), CG (which contains the median CS), and MK have a common point, which is S . Thus Q lies on midline MS .

The same reasoning proves that P lies on midline MR . Since all homothety ratios are the same, $\frac{MQ}{MS} = \frac{MP}{MR}$, which shows that MPQ is similar to MRS , which in turn is similar to ABC , and we are done.

Problem 2

Suppose there are 997 points given in a plane. If every two points are joined by a line segment with its midpoint coloured in red, show that there are at least 1991 red points in the plane. Can you find a special case with exactly 1991 red points?

Solution

Embed the points in the cartesian plane such that no two points have the same y -coordinate. Let P_1, P_2, \dots, P_{997} be the points and $y_1 < y_2 < \dots < y_{997}$ be their respective y -coordinates. Then the y -coordinate of the midpoint of $P_i P_{i+1}$, $i = 1, 2, \dots, 996$ is $\frac{y_i + y_{i+1}}{2}$ and the y -coordinate of the midpoint of $P_i P_{i+2}$, $i = 1, 2, \dots, 995$ is $\frac{y_i + y_{i+2}}{2}$. Since

$$\frac{y_1 + y_2}{2} < \frac{y_1 + y_3}{2} < \frac{y_2 + y_3}{2} < \frac{y_2 + y_4}{2} < \dots < \frac{y_{995} + y_{997}}{2} < \frac{y_{996} + y_{997}}{2},$$

there are at least $996 + 995 = 1991$ distinct midpoints, and therefore at least 1991 red points. The equality case happens if we take $P_i = (0, 2i)$, $i = 1, 2, \dots, 997$. The midpoints are $(0, i + j)$, $1 \leq i < j \leq 997$, which are the points $(0, k)$ with $1 + 2 = 3 \leq k \leq 996 + 997 = 1993$, a total of $1993 - 3 + 1 = 1991$ red points.

Problem 3

Let $a_1, a_2, \dots, a_n, b_1, b_2, \dots, b_n$ be positive real numbers such that $a_1 + a_2 + \dots + a_n = b_1 + b_2 + \dots + b_n$. Show that

$$\frac{a_1^2}{a_1 + b_1} + \frac{a_2^2}{a_2 + b_2} + \dots + \frac{a_n^2}{a_n + b_n} \geq \frac{a_1 + a_2 + \dots + a_n}{2}.$$

Solution

By the Cauchy-Schwartz inequality,

$$\left(\frac{a_1^2}{a_1 + b_1} + \frac{a_2^2}{a_2 + b_2} + \dots + \frac{a_n^2}{a_n + b_n} \right) ((a_1 + b_1) + (a_2 + b_2) + \dots + (a_n + b_n)) \geq (a_1 + a_2 + \dots + a_n)^2.$$

Since $((a_1 + b_1) + (a_2 + b_2) + \dots + (a_n + b_n)) = 2(a_1 + a_2 + \dots + a_n)$,

$$\frac{a_1^2}{a_1 + b_1} + \frac{a_2^2}{a_2 + b_2} + \dots + \frac{a_n^2}{a_n + b_n} \geq \frac{(a_1 + a_2 + \dots + a_n)^2}{2(a_1 + a_2 + \dots + a_n)} = \frac{a_1 + a_2 + \dots + a_n}{2}.$$

Problem 4

During a break, n children at school sit in a circle around their teacher to play a game. The teacher walks clockwise close to the children and hands out candies to some of them according to the following rule. He selects one child and gives him a candy, then he skips the next child and gives a candy to the next one, then he skips 2 and gives a candy to the next one, then he skips 3, and soon. Determine the values of n for which eventually, perhaps after many rounds, all children will have at least one candy each.

Answer: All powers of 2.

Solution 1

Number the children from 0 to $n - 1$. Then the teacher hands candy to children in positions $f(x) = 1 + 2 + \dots + x \pmod n = \frac{x(x+1)}{2} \pmod n$. Our task is to find the range of $f: \mathbb{Z}_n \rightarrow \mathbb{Z}_n$, and to verify whether the range is \mathbb{Z}_n , that is, whether f is a bijection.

If $n = 2^a m$, $m > 1$ odd, look at $f(x)$ modulo m . Since m is odd, $m \mid f(x) \iff m \mid x(x+1)$. Then, for instance, $f(x) \equiv 0 \pmod m$ for $x = 0$ and $x = m - 1$. This means that $f(x)$ is not a bijection modulo m , and there exists t such that $f(x) \not\equiv t \pmod m$ for all x . By the Chinese Remainder Theorem,

$$f(x) \equiv t \pmod n \iff \begin{cases} f(x) \equiv t \pmod{2^a} \\ f(x) \equiv t \pmod m \end{cases}$$

Therefore, f is not a bijection modulo n .

If $n = 2^a$, then

$$f(x) - f(y) = \frac{1}{2}(x(x+1) - y(y+1)) = \frac{1}{2}(x^2 - y^2 + x - y) = \frac{(x-y)(x+y+1)}{2}.$$

and

$$f(x) \equiv f(y) \pmod{2^a} \iff (x-y)(x+y+1) \equiv 0 \pmod{2^{a+1}}. \quad (*)$$

If x and y have the same parity, $x+y+1$ is odd and $(*)$ is equivalent to $x \equiv y \pmod{2^{a+1}}$. If x and y have different parity,

$$(*) \iff x+y+1 \equiv 0 \pmod{2^{a+1}}.$$

However, $1 \leq x+y+1 \leq 2(2^a - 1) + 1 = 2^{a+1} - 1$, so $x+y+1$ is not a multiple of 2^{a+1} . Therefore f is a bijection if n is a power of 2.

Solution 2

We give a full description of a_n , the size of the range of f .

Since congruences modulo n are defined, via Chinese Remainder Theorem, by congruences modulo p^α for all prime divisors p of n and α being the number of factors p in the factorization of n , $a_n = \prod_{p^\alpha \parallel n} a_{p^\alpha}$.

Refer to the first solution to check the case $p = 2$: $a_{2^\alpha} = 2^\alpha$.

For an odd prime p ,

$$f(x) = \frac{x(x+1)}{2} = \frac{(2x+1)^2 - 1}{8},$$

and since p is odd, there is a bijection between the range of f and the quadratic residues modulo p^α , namely $t \mapsto 8t + 1$. So a_{p^α} is the number of quadratic residues modulo p^α .

Let g be a primitive root of p^α . Then there are $\frac{1}{2}\phi(p^\alpha) = \frac{p-1}{2} \cdot p^{\alpha-1}$ quadratic residues that are coprime with p : $1, g^2, g^4, \dots, g^{\phi(p^\alpha)-2}$. If p divides a quadratic residue kp , that is, $x^2 \equiv kp \pmod{p^\alpha}$, $\alpha \geq 2$, then p divides x and, therefore, also k . Hence p^2 divides this quadratic residue, and these quadratic residues are p^2 times each quadratic residue of $p^{\alpha-2}$. Thus

$$a_{p^\alpha} = \frac{p-1}{2} \cdot p^{\alpha-1} + a_{p^{\alpha-2}}.$$

Since $a_p = \frac{p-1}{2} + 1$ and $a_{p^2} = \frac{p-1}{2} \cdot p + 1$, telescoping yields

$$a_{p^{2t}} = \frac{p-1}{2}(p^{2t-1} + p^{2t-3} + \cdots + p) + 1 = \frac{p(p^{2t} - 1)}{2(p+1)} + 1$$

and

$$a_{p^{2t-1}} = \frac{p-1}{2}(p^{2t-2} + p^{2t-4} + \cdots + 1) + 1 = \frac{p^{2t} - 1}{2(p+1)} + 1$$

Now the problem is immediate: if n is divisible by an odd prime p , $a_{p^\alpha} < p^\alpha$ for all α , and since $a_t \leq t$ for all t , $a_n < n$.

Problem 5


Given are two tangent circles and a point P on their common tangent perpendicular to the lines joining their centres. Construct with ruler and compass all the circles that are tangent to these two circles and pass through the point P .

Solution

Throughout this problem, we will assume that the given circles are *externally* tangent, since the problem does not have a solution otherwise.

Let Γ_1 and Γ_2 be the given circles and T be their tangency point. Suppose ω is a circle that is tangent to Γ_1 and Γ_2 and passes through P .

Now invert about point P , with radius PT . Let any line through P that cuts Γ_1 do so at points X and Y . The power of P with respect to Γ_1 is $PT^2 = PX \cdot PY$, so X and Y are swapped by this inversion. Therefore Γ_1 is mapped to itself in this inversion. The same applies to Γ_2 . Since circle ω passes through P , it is mapped to a line tangent to the images of Γ_1 (itself) and Γ_2 (also itself), that is, a common tangent line. This common tangent cannot be PT , as PT is also mapped to itself. Since Γ_1 and Γ_2 have exactly other two common tangent lines, there are two solutions: the inverses of the tangent lines.



We proceed with the construction with the aid of some macro constructions that will be detailed later.

Step 1. Draw the common tangents to Γ_1 and Γ_2 .

Step 2. For each common tangent t , draw the projection P_t of P onto t .

Step 3. Find the inverse P_1 of P_t with respect to the circle with center P and radius PT .


Step 4. ω_t is the circle with diameter PP_1 .

Let's work out the details for steps 1 and 3. Steps 2 and 4 are immediate.

Step 1. In this particular case in which Γ_1 and Γ_2 are externally tangent, there is a small shortcut:

- Draw the circle with diameter on the two centers O_1 of Γ_1 and O_2 of Γ_2 , and find its center O .

- Let this circle meet common tangent line OP at points Q, R . The required lines are the perpendicular to OQ at Q and the perpendicular to OR at R .



Let's show why this construction works. Let R_i be the radius of circle Γ_i and suppose without loss of generality that $R_1 \leq R_2$. Note that $OQ = \frac{1}{2}O_1O_2 = \frac{R_1+R_2}{2}$, $OT = OO_1 - R_1 = \frac{R_2-R_1}{2}$, so

$$\sin \angle TQO = \frac{OT}{OQ} = \frac{R_2 - R_1}{R_1 + R_2},$$

which is also the sine of the angle between O_1O_2 and the common tangent lines.


Let t be the perpendicular to OQ through Q . Then $\angle(t, O_1O_2) = \angle(OQ, QT) = \angle TQO$, and t is parallel to a common tangent line. Since

$$d(O, t) = OQ = \frac{R_1 + R_2}{2} = \frac{d(O_1, t) + d(O_2, t)}{2},$$


and O is the midpoint of O_1O_2 , O is also at the same distance from t and the common tangent line, so these two lines coincide.

Step 3. Finding the inverse of a point X given the inversion circle Ω with center O is a well known procedure, but we describe it here for the sake of completeness.

- If X lies in Ω , then its inverse is X .
- If X lies in the interior of Ω , draw ray OX , then the perpendicular line ℓ to OX at X . Let ℓ meet Ω at a point Y . The inverse of X is the intersection X' of OX and the line perpendicular to OY at Y . This is because OYX' is a right triangle with altitude YX , and therefore $OX \cdot OX' = OY^2$.
- If X is in the exterior of Ω , draw ray OX and one of the tangent lines ℓ from X to Ω (just connect X to one of the intersections of Ω and the circle with diameter OX). Let ℓ touch Ω at a point Y . The inverse of X is the projection X' of Y onto OX . This is because OYX' is a right triangle with altitude YX' , and therefore $OX \cdot OX' = OY^2$.



X is inside Ω



X is outside Ω